

# The Role of Candidate-Gene *CNTNAP2* in Childhood Apraxia of Speech and Specific Language Impairment

TM Centanni,<sup>1,2</sup> JN Sanmann,<sup>3</sup> JR Green,<sup>1</sup> J Iuzzini-Seigel,<sup>1,4</sup> C Bartlett,<sup>5</sup> WG Sanger,<sup>3†</sup> and TP Hogan<sup>1\*</sup>

<sup>1</sup>MGH Institute of Health Professions, Boston, Massachusetts

<sup>2</sup>Massachusetts Institute of Technology, Cambridge, Massachusetts

<sup>3</sup>University of Nebraska Medical Center, Nebraska Medical Center, Omaha, Nebraska

<sup>4</sup>Marquette University, Milwaukee, MI

<sup>5</sup>The Ohio State University, Columbus, Ohio

Manuscript Received: 17 December 2014; Manuscript Accepted: 13 May 2015

Childhood apraxia of speech (CAS) is a debilitating pediatric speech disorder characterized by varying symptom profiles, comorbid deficits, and limited response to intervention. Specific Language Impairment (SLI) is an inherited pediatric language disorder characterized by delayed and/or disordered oral language skills including impaired semantics, syntax, and discourse. To date, the genes associated with CAS and SLI are not fully characterized. In the current study, we evaluated behavioral and genetic profiles of seven children with CAS and eight children with SLI, while ensuring all children were free of comorbid impairments. Deletions within *CNTNAP2* were found in two children with CAS but not in any of the children with SLI. These children exhibited average to high performance on language and word reading assessments in spite of poor articulation scores. These findings suggest that genetic variation within *CNTNAP2* may be related to speech production deficits. © 2015 Wiley Periodicals, Inc.

**Key words:** gene variant; speech production; CAS; SLI

## INTRODUCTION

Childhood apraxia of speech (CAS) is a debilitating pediatric speech disorder that affects 1–2 children per thousand [Shriberg, Aram, & Kwiatkowski, 1997] and is often resistant to intervention [Lewis, Freebairn, Hansen, Iyengar, & Taylor, 2004; Teverovsky, Bickel, & Feldman, 2009]. CAS is characterized by a broad range of speech abnormalities that affect accuracy and consistency of speech sound production, as well as suprasegmental features such as prosody [Marquardt, 2004; Iuzzini, 2012], or intonation [American Speech-Language-Hearing, 2007]. Children with CAS may evidence concomitant deficits such as language impairment, dysarthria, or intellectual disability [American Speech-Language-Hearing, 2007]. Comorbid deficits contribute to the complex

### How to Cite this Article:

Centanni TM, Sanmann JN, Green JR, Iuzzini-Seigel J, Bartlett C, Sanger WG, Hogan TP. 2015. The Role of Candidate-Gene *CNTNAP2* in Childhood Apraxia of Speech and Specific Language Impairment. *Am J Med Genet Part B* 9999:1–8.

speech presentation observed in these children, which has led to challenges with early diagnosis and has motivated the search for reliable genetic markers. Specific Language Impairment (SLI) is a persistent developmental language impairment that is characterized by delayed and/or disordered oral language skills, including impaired discourse, syntax, and semantics. The high comorbidity between CAS and SLI [Shriberg, Tomblin, & McSweeney, 1999] makes it extremely difficult to determine which genes are related to the phenotypes of CAS and which are related to SLI, as opposed to which are associated with comorbid CAS-LI.

As part of a larger study on speech and language impairments, we evaluated the genetic profiles of children with a diagnosis of CAS or SLI. This sample was strictly controlled so that no children in the

<sup>†</sup>In memory of our colleague and collaborator, Warren G Sanger, who passed away during the preparation of this manuscript. The authors have no conflicts of interest to disclose.

Grant sponsor: University of Nebraska Health Research Consortium; Grant sponsor: University of Nebraska-Lincoln Barkley Memorial Trust.

\*Correspondence to:

Tiffany P. Hogan, 36 1st Ave, Boston, MA 02129.

E-mail: thogan@mghihp.edu

Article first published online in Wiley Online Library (wileyonlinelibrary.com): 00 Month 2015

DOI 10.1002/ajmg.b.32325

CAS group had comorbid reading, language, or cognitive impairments and no children in the SLI group had comorbid cognitive or articulation impairments. In this report, we discuss the speech and language phenotypes of two children with CAS who had deletions in the region of chromosome 7 that contains the neurexin gene *CNTNAP2* (7q35). *CNTNAP2* is located downstream from and is regulated by *FOXP2* (7q31), which has been linked to the occurrence of CAS in the oft-studied KE family [Lai, Fisher, Hurst, Vargha-Khadem, & Monaco, 2001; Vernes et al., 2008]. *CNTNAP2* is closely related to *FOXP2* and has been identified as a candidate gene for dyslexia, SLI, and autism [Laffin et al., 2012; Rodenas-Cuadrado, Ho, & Vernes, 2014]. To our knowledge, this is the first report to link variants in *CNTNAP2* to CAS without comorbid reading, language, and cognitive impairments, which indicates that *CNTNAP2* variants may be associated with deficits in speech production in the absence of comorbid reading, language, and cognitive impairments.

MATERIALS AND METHODS

Participants

Fifteen children ranging in age from 4;5–17;2 (years; months) participated as part of a larger study on the biological pathways of speech and language disorders. All procedures were approved by the Institutional Review Board of the University of Nebraska Medical Center and the University of Nebraska-Lincoln, and all participants were consented prior to participation. Participants underwent a series of commonly administered, age-appropriate speech, language, reading, and cognitive assessments including the *Goldman Frisloe Test of Articulation-2nd Edition* [GFTA-2; Goldman & Frisloe, 2000], the *Clinical Evaluation of Language Fundamentals-Fourth Edition* [CELF-4; Semel et al., 2003], *Reynolds Intellectual Assessment Scales* [RIAS; Reynolds & Kamphaus, 2003], and the *Woodcock Reading Mastery Test-Revised* [WRMT-R; Woodcock, 1998]. All participants were required to have normal cognition based on a standard score higher of 75 or higher on the RIAS.

Group Assignment

Participants in the CAS group were typically referred to the study with a history of CAS diagnosis and treatment by a clinician with expertise in CAS. The CAS diagnosis was confirmed if the participant evidenced at least 4 of 11 features associated with CAS [adapted from Shriberg, Potter, & Strand, 2011] during a standardized, norm-referenced articulation assessment [Goldman & Frisloe, 2000]. We reasoned that if a child with a history of CAS—who did not have comorbid language impairment, cognitive defi-

cit, or dysarthria—produced a high number of features on simple test items, we could be more certain in confirming the CAS diagnosis rather than a different deficit (e.g., dyslexia), which could yield numerous errors on complex items [Catts, 1989]. Therefore, two trained raters, a speech language pathologist with expertise in CAS and a speech language pathology graduate student, independently blind-rated each child’s responses on the GFTA-2 using the operational definitions in Table III.

Participants in this group were also required to have an articulation test (GFTA-2) percentile score at or below the 16th percentile and a normal language standard score of 85 or higher on the CELF-4. None of the participants included in the CAS group in this study reported a history of diagnosis or treatment for language impairment.

Children were assigned to the SLI group based on GFTA-2 percentile scores of 16 or higher, fewer than 4/11 CAS features, and a CELF-4 standard score below 85. See Table I for inclusion criterion and Table II for assessment scores.

DNA Collection and Isolation

Buccal cell samples were collected from participants using the Isohelix DNA swab packs (Cell Projects, Ltd., Kent, United Kingdom), and DNA was extracted per manufacturer’s recommendations using the QIAcube (Qiagen, Valencia, CA). DNA quantity and quality were determined using the NanoDrop ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE) and agarose gel electrophoresis, respectively.

High-Resolution Genome-Wide Analysis

High-resolution genome-wide analysis was performed on genomic DNA using the CytoScanHD™ array (Affymetrix, Santa Clara, CA) according to manufacturer’s instruction. This array contains more than 2.6 million markers for high-resolution whole-genome copy number analysis and 750,000 genotype-able single nucleotide polymorphisms (SNPs) for reliable detection of copy neutral loss of heterozygosity (CN-LOH). Data were visualized and analyzed with the Chromosome Analysis Suite (ChAS) software (Affymetrix) using the following filter parameters: (1) ≥25 markers and ≥5 kilobases (kb) for copy number variants (CNVs) and (2) ≥5 megabases (Mb) for CN-LOH.

Reliability of Perceptual Feature Ratings

To ensure confidence in group assignments inter-rater reliability of perceptual feature ratings was calculated on data from all participants. The intra-rater correlation coefficient with absolute error in

TABLE I. Inclusion Criterion for Both Groups

	Nonverbal IQ (standard score)	Speech production (percentile)	Language (standard score)	Word reading (standard score)	# of CAS features
CAS (N = 7)	>75	≤16th percentile	≥85	Any	≥4
SLI (N = 8)	>75	>16th percentile	<85	Any	<4

TABLE II. Behavioral Profiles for Both Groups. \*\*\* $p < 0.001$ 

	Nonverbal IQ [standard score]	Speech production [percentile]***	Language [standard score]***	Word reading [standard score]	# of CAS features***
CAS (N = 7)	110.43 [11.60]	3.14 [5.24] -percentile	109.57 [16.35]	101.57 [7.70]	7.14 [2.19]
SLI (N = 8)	102.33 [4.95]	34.44 [6.42] - percentile	73.67 [9.77]	97.89 [9.92]	1.44 [1.01]

parenthesis was .98 (.30 features), showing a high level of agreement between raters for perceptual feature rating using the operational definitions that are included in Table III.

## RESULTS

### CNTNAP2 Deletions in Children With CAS

In the current study, 2 of the 7 children with CAS exhibited deletions within the *CNTNAP2* gene, at 7q35, with array probes encompassing 6.77 kilo-bases (kb) in length. Deletion breakpoints were identical in both children spanning 147714709–147721486 bp (human GRCh37/hg19 assembly). The deletions were located in alternative intron 18 and were approximately 20 kb away from the

nearest exon (Fig. 1). Unfortunately, we were unable to collect genetic samples from the participants' biological parents, so we could not determine if these deletions were inherited or de novo in nature. Child 1 was a 12-year-old female and child 2 was an 8-year-old male (see Table IV for assessment scores).

### Behavioral Profiles of Children With CNTNAP2 Deletions

**Nonverbal IQ.** Both participants evidenced normal nonverbal intelligence. Child 1 exhibited average intelligence ( $SS = 95$ ) and child 2 exhibited high intelligence ( $SS = 131$ ), more than two standard deviations above the mean (Table IV).

TABLE III. Operational Definitions for CAS Characteristics

Characteristic	Definition
Vowel error	A vowel production error in which the vowel is substituted for another phoneme OR in which the vowel is recognizable as a specific phoneme but it is not produced exactly correctly (e.g., not a prototypical production, may sound like it's in between two vowels). It is not considered an error if the vowel is substituted with another phoneme that is consistent with an adult-like model (e.g., /hat dag/ /hat dog/)
Consonant distortion	A consonant production error in which a speech sound is recognizable as a specific phoneme but it is not produced exactly correctly (e.g., an /s/ that is produced with lateralization or dentalization).
Stress errors	An error in which the appropriate stress is not produced correctly. For example: conDUCT vs. CONduct have different stress patterns. It is considered an error if the stress is inappropriately equalized across syllables, or on the wrong syllable.
Syllable segregation	Brief or lengthy pause between syllables which is not appropriate.
Groping; prevocalic	[silent] articulatory searching prior to onset of phonation, possibly in an effort to improve the accuracy of the production. Video is needed to assess this feature.
Intrusive schwa (e.g. in clusters)	A schwa is added in between consonants. For example, it may be inserted in between the consonants in a cluster (e.g., /blu/ becomes /bəl u/). This NOT considered a "vowel error"
Voicing errors	A sound is produced as its voicing cognate (e.g., a /p/ that is produced as a /b/). In addition, this could also describe productions which appear to be in between voicing categories (e.g., blurring of voicing boundaries).
Slow rate	Speech rate is not typical. It is slower during production of part (e.g., zzziiiiiper/zipper) or the whole word (e.g., tooommmmaaatoooo/tomato).
Increased difficulty with multisyllabic words	The participant has a disproportionately increased number of errors as the number of syllables increases (as compared to words with fewer syllables).
Resonance or nasality disturbance	Sounds either <i>hyponasal</i> : not enough airflow out of nose/"stuffy" OR <i>hypernasal</i> : too much airflow out of nose for non-nasal phonemes (e.g., plosives).
Difficulty achieving initial articulatory configurations or transitional movement gestures	Initiation of utterance or initial speech sound may be difficult for child to produce and may sound lengthened or uncoordinated. Also, child may evidence lengthened or disrupted coarticulatory gestures or movement transitions from one sound to the next.




FIG. 1. Location of the deletion reported here within *CNTNAP2* in the context of *CNTNAP2*'s location on chromosome 7. The deletion reported here was located in alternative intron 18 and was 20 kb from the nearest exon. Images were adapted from the UCSC genome database (<https://genome.ucsc.edu/>).

**CAS features.** The number of CAS features differed between the cases (Table IV). Child 1 had four features where participant 2 had nine features (out of a possible 11). The participants evidenced four CAS features in common, which included: vowel errors, consonant distortions, excessive or equal stress, and voicing errors. Participant 2 also evidenced syllable segregation, groping, slow rate, resonance disturbance, and difficulty achieving initial articulatory configuration or transitional movement gestures.

**Speech severity.** Both children scored in the 1st percentile on the speech production assessment. This indicates that both children evidenced severe speech deficits relative to their same aged peers. (Table IV)

**Language and literacy.** Both children evidenced normal language and word reading ability. On the *CELF-4* core language test, child 1 had a standard score of 121 and child 2 a score of 111 indicating normal-to-high language ability for these participants (Table III). On the *WRMT* reading assessment, child 1 had a standard score of 105 and child 2 a score of 110 (Table IV). These results suggest that, in some children with CAS, deletions containing *CNTNAP2* may play a role in speech production without impacting language skills or reading ability.

Although it is possible that at the time of testing our participants evidenced remediated language impairments, their performance in the normal-to-high range on language and reading tests suggest that a history of language impairment is unlikely. Longitudinal research on children with language impairment [Stothard, Snowling, Bishop, Chipchase, & Kaplan, 1998] shows that children with a

preschool history of language impairment who were retested during adolescence and scored in the normal range, still evidenced difficulty with phonological processing and literacy skills. Given that neither family reported difficulty with language or reading impairments, nor did either participant demonstrate performance at the lower end of normal on language or reading assessments, we feel confident that these children likely had CAS speech symptoms in the absence of language impairment.

Additional Genetics Findings

We note that both children had two additional CNVs that met report5 criteria (Table V). Child 1 had deletions at 5q34 and 12p12.3. Previous studies have associated CNVs at 5q34 with facial abnormalities such as cleft lip, depressed nasal bridge, and microcephaly [Schafer et al., 2001; Chen et al., 2012]. Previous associations in this region included the genes *MSX2*, *NKX2-5*, and *NSD1*, none of which were included in the CNV seen in Child 1 (deletion was small and included only *ODZ2*). CNVs at 12p12.3 have been previously associated with language delay, dysmorphic features, and hypotonia [Gläser et al., 2003; Lamb et al., 2012]. These previous studies focused on nearby regions and the gene *SOX5*, which was not affected by the deletion seen in Child 1.

Child 2 had deletions at 1q25.1 and 15q21.2. Deletion at 1q25.1 has previously been linked to delayed language and expressive language impairment [Höglund, Jalkanen, Marttinen, & Alitalo, 2003], though the deletion previously reported was significantly

TABLE IV. Assessment Scores for Two Children With *CNTNAP2* Deletions and CAS

	Nonverbal IQ (standard score)	Speech production (percentile)	Language (standard score)	Word reading (standard score)	# of CAS features
Child 1	95	1	121	105	4/11
Child 2	131	1	111	110	9/11

TABLE V. Summary of Additional CNVs in Two Children With CAS

	CNV identified	Type	Size (kbp)	Linear location	Previous associations
Child 1	5q34	Deletion	26.052	167061974- 167088026	cleft lip, depressed nasal bridge, microcephaly [Chen et al., 2012; Schafer et al., 2001]
	12p12.3	Deletion	47.118	15826648- 15873766	language delay, dysmorphic features and hypotonia [Gläser et al., 2003; Lamb et al., 2012]
Child 2	1q25.1	Deletion	224.684	174450384- 174675068	delayed language and expressive language impairment [Höglund et al., 2003]
	15q21.2	Deletion	24.483	51800269 - 51824752	abnormal facial shape and muscular hypotonia [Moreno-De-Luca et al., 2011]

larger than the one seen in Child 2. Deletion at 15q21.2 has previously been associated with abnormal facial shape and muscular hypotonia [Moreno-De-Luca et al., 2011]. This previous deletion did not overlap with the deletion seen in Child 2. The differences between the deletions seen in our cases and CNVs reported in previous literature linking these regions to speech and language delay support the hypothesis that the deletion seen in the *CNTNAP2* gene in the current study may have a functional consequence on phenotype. Formal evaluation by a board-certified clinical geneticist was not performed for either participant. Additional studies will be needed to determine whether CNVs in these regions contribute to speech production deficits.

### Lack of *CNTNAP2* Variants in Children With SLI

Although *CNTNAP2* variants have predominantly been linked to the presence of SLI, we did not observe any CNVs at 7q35 in our sample of eight children with this disorder. The clinical significance of the CNVs identified in the children with SLI in other areas of the genome are currently being analyzed for a future report.

### In Silico Analysis of the Deleted Region

Since the deletion is intronic, the putative mechanism for how this CNV could affect cellular function does not involve protein sequence changes. We examined public data sources for evidence that this region has regulatory potential [Karolchik et al., 2014]. Maunakea et al. [2010] showed two methylation peaks and two RNA-seq peaks within 2.2 kb in an adult human brain. There are two distinct 5' SMART tags indicating two transcription start sites in opposite directions. These data are indicative of an enhancer RNA. Corroborating these data, the ENCODE database [Kellis et al., 2014] shows a DNaseI hypersensitive site in four cell lines (H9es, H1-hESC, lps, NT2-D1) and, overlapping this DNaseI hypersensitive site, ChIP-seq indicates binding of twelve transcription factors (NANOG, MAFK, CEBPB, EP300, BCL11A, JUND, TEAD4, SP1, POU5F1, TCF12, SIN3A, HDAC2). Taken together, these data indicate that a bidirectional enhancer RNA is located within the 6.7 kb deletion. However, it is not possible to infer if the enhancer RNA acts in cis to affect *CNTNAP2* expression or in trans to regulate a different gene.

## DISCUSSION

### Summary of Results

In the current study, we describe two children with a strictly controlled diagnosis of childhood apraxia of speech (CAS) without comorbid reading, language, or cognitive impairments who had deletions in alternative intron 18 of *CNTNAP2* at 7q35. Both children with CAS evidenced normal language and word reading abilities. In contrast, we did not observe any variants in *CNTNAP2* in the eight children with SLI. These results suggest that in some children, deletions containing *CNTNAP2* may play a role in speech production in the presence of intact language-based processing skills.

### Evaluation of Study Design

Strengths of the current study include absence of comorbid language and reading deficits and the precise speech, language, and reading phenotyping of each child, including the strict diagnostic criteria used to define CAS. In prior research, deletions containing *CNTNAP2* have been associated with a variety of diagnoses including autism, language impairment, speech delay, and dyslexia [Peter et al., 2011; Laffin et al., 2012]. The interpretation of these studies, however, is challenged by the high likelihood of comorbid speech motor problems in these cohorts. Our result that *CNTNAP2* variants occurred in multiple children with CAS in the absence of reading, language, and cognitive impairments raises the possibility that *CNTNAP2* variants are involved in the motor components of speech. One caveat of this study is that our sample size is small; therefore, we are cautious in making any definitive conclusions about these CNVs being causative for CAS.

### Intron Versus Exon Deletion

Though the deletions reported here did not contain any exons, previous work in other disorders report intronic deletions in *CNTNAP2* that were associated with disorder phenotypes. For example, a recent paper described the case of a woman with epilepsy and schizophrenia who also had a small intron deletion in *CNTNAP2*, while her twin sister was free of both disorders and did not have this deletion [Friedman, Vrijenhoek, & Markx, 2007]. Other studies have linked intronic deletions in *CNTNAP2* to a



variety of features, including age of first word, receptive and expressive language impairment, and non-word repetition [Poot, Beyer, Schwaab, & Damatova, 2010; Peñagarikano & Geschwind, 2012]. Though larger sample sizes and functional studies are needed to more decisively determine the role of intronic CNVs on phenotype, our work supports previous reports that intronic deletions in *CNTNAP2* may have functional consequences on speech production abilities.

### Prevalence of *CNTNAP2* CNV

Though the sample size in the current study is small, recent work suggests the presence of this specific CNV in 2 of our 7 children with CAS may deviate from expectations relative to the general population. The prevalence of this CNV in our sample is 14.3%. To compare with two population-based samples, we considered only CNVs that overlap with the one described here, yet are still within the same intron (i.e., no exonic involvement). The first study found this variant in 0.349% of a Swiss European sample [N = 717; Vogler et al., 2010] and the second study found this variant in 1.74% of an Ontario population sample [30/873; Costain et al., 2013]. The difference in minor allele frequency is statistically significant across the Vogler and Costain studies (Fisher's exact test;  $P < 0.001$ ) that could be due to a difference in probe density across array platforms. The Vogler study used the Affymetrix Genome-Wide Human SNP Array 6.0 while the Ontario study was CytoHD specific—the same technology as in our study. The difference could also be due to the fact that these populations were not strictly controlled for speech and language skills. The prevalence between our population and the Ontario study is different (28% vs. 1.74%; Fisher's exact test,  $P = 0.024$ ), and in concert with the prevalence estimates of the disorder in the population [Shriberg, Aram, & Kwiatkowski, 1997], suggest that this variant is more likely to occur in individuals with apraxia than by chance in the general population. Additional studies are needed to confirm that this CNV is more prevalent in children with CAS than a verified typically developing control population.

### Biological Mechanisms of *CNTNAP2* Variants

The downstream effect of this intronic CNV has not been developed in the literature. However, several studies define typical *CNTNAP2* regulation, and more recently, also define the effects of genetic variation on *CNTNAP2* functioning in humans [Zeeland, 2010; Dennis & Jahanshad, 2011; Hohenberg & Wigand, 2013]. *CNTNAP2* is a cell-cell adhesion molecular and member of the neuroligin protein family [Poliak, Gollan, Martinez, & Custer, 1999]. *CNTNAP2* mediates neuron-glia interactions at juxtaparanodal axonal regions and from that role is involved in myelination [Poliak & Gollan, 2001]. During neuronal differentiation, *CNTNAP2* also localizes potassium channels giving it a dual role for propagation of action potentials within the juxtaparanodal region [Rasband, 2004]. Given this association with myelination and functioning of axons, studies have examined white matter in connection with *CNTNAP2* variation using in vivo tractography techniques. Thus far, these studies do observe significant difference between typically developing individuals carrying autism or

language-impairment associated variants versus persons without those variants [Zeeland, 2010; Dennis & Jahanshad, 2011], and more recently a variant not previously associated with disease has also been found [Hohenberg & Wigand, 2013]. Interpreting these in vivo tractography measures in the context of neurobiological function is more challenging. Additionally, it is clear that *CNTNAP2* is involved in other processes that will require additional study designs to fully elucidate. An example includes studying epigenetic regulation, where differences between humans and chimpanzees has been demonstrated [Schneider, Hajj, & Richter, 2014].

### The Role of *CNTNAP2* in Motor Impairments

Since the speech production-related gene *FOXP2* regulates *CNTNAP2*, one unexplored possibility is that the observed association between *CNTNAP2* and disorders such as dyslexia or autism is due to comorbid speech motor disturbances rather than to a reading or language impairment alone. Recent studies in animal models support this hypothesis. *CNTNAP2* [Condro & White, 2014] is important for song mimicking in birds, which is a process that is highly dependent on motor control. In addition, recent studies in individuals with dyslexia have reported associations between sequence variations in *CNTNAP2* and stuttering [Petrin et al., 2010] as well as difficulties with motor-heavy tasks such as rapid oral reading [Peter et al., 2011]. In addition, a recent study also reported an intron deletion in *CNTNAP2* in a child with CAS and a small insertion in two other children with the disorder [Worthey et al., 2013]. These findings support our finding that abnormalities in *CNTNAP2* can occur in persons with CAS without comorbid reading and language impairment. Finding *CNTNAP2* variants in children with CAS who do not have a language or reading impairment would indicate that *CNTNAP2* may adversely affect the pathway responsible for speech production, rather than the pathway responsible for language and reading, in these children.

### ACKNOWLEDGMENTS

This research was supported by the University of Nebraska Health Research Consortium (Co-PIs Green and Hogan) and the University of Nebraska-Lincoln Barkley Memorial Trust. The authors wish to thank Shelley Smith for guidance during experimental design. The authors also wish to thank Kimber Green, Sara Benham, Dyann Rupp, Tacy Corson, Phoebe Chung, Natalie Covington, and Kristin Schneller for their assistance with data collection and Diane Pickering and Danielle Bishay for specimen processing and genetic data analysis.

### REFERENCES

- American Speech-Language-Hearing 2007. Childhood apraxia of speech. [Technical Report]. Available from [www.asha.org/policy](http://www.asha.org/policy).
- Catts H. 1989. Speech production deficits in developmental dyslexia. *J Speech Hear Disord* 54(3):422–428.
- Chen C, Lin S, Chen M, Y Su, S Chern, Liu Y, Wang W. 2012. Partial monosomy 3p (3p26.2->pter) and partial trisomy 5q (5q34->qter) in a

- girl with coarctation of the aorta, congenital heart defects, short stature, microcephaly and developmental delay. *Genet Counsel* 23(3):405–413.
- Condro MC, White SA. 2014. Recent advances in the genetics of vocal learning. *Comp Cogn Behav Rev* 9.
- Costain G, Lionel AC, Merico D, Forsythe P, Russel K, Lowther C, Yuen T, Husted J, Stavropoulos DJ, Speevak M, Chow EW, Marshall CR, Scherer SW, Bassett AS. 2013. Pathogenic rare copy number variants in community-based schizophrenia suggest a potential role for clinical microarrays. *Human Molecular Genetics* 22:4485–4501.
- Dennis E, Jahanshad N. 2011. Altered structural brain connectivity in healthy carriers of the autism risk gene *CNTNAP2*. *Brain* 1(6): 447–459.
- Friedman J, Vrijenhoek T, Markx S. 2007. *CNTNAP2* gene dosage variation is associated with schizophrenia and epilepsy. *Mol Psychiatr* 13(3):261–266.
- Gläser B, Rossier E, Barbi G, Chiaie L D, Blank C, Vogel W, Kehrer-Sawatzki H. 2003. Molecular cytogenetic analysis of a constitutional de novo interstitial deletion of chromosome 12p in a boy with developmental delay and congenital anomalies. *Am J Med Genet Part A* 116A(1): 66–70. doi: 10.1002/ajmg.a.10878
- Goldman R, Fristoe M. 2000. Goldman-Fristoe test of articulation-2 (GFTA-2). Circle Pines, MN: American Guidance Service.
- Höglund P, Jalkanen R, Marttinen E, Alitalo T. 2003. Interstitial 1q25.3-q31.3 deletion in a boy with mild manifestations. *Am J Med Genet Part A* 123A(3):290–295. doi: 10.1002/ajmg.a.20385
- Hohenberg C, von, Wigand M. 2013. *CNTNAP2* polymorphisms and structural brain connectivity: A diffusion-tensor imaging study. *J Psychiatr* 47(10):1349–1356.
- Iuzzini J. 2012. Inconsistency of speech in children with childhood apraxia of speech, phonological disorders, and typical speech. *Vasa* 73(07).
- Karolchik D, Barber GP, Casper J, Clawson H, Cline MS, Diekhans M, Dreszer et al. 2014. The UCSC genome browser database: 2014 update. *Nucleic Acids Research* 42(D1):D764–D770.
- Kellis M, Wold B, Snyder MP, Berstein BE, Kundaje A, Marinov GK, Ward LD, Birney E, Crawford GE, Dekker J, Dunham I, Elnitski LL, Farnham PJ, Feingold EA, Gerstein M, Giddings MC, Gilbert DM, Gingeras TR, Greed ED, Guigo R, Hubbard T, Kent J, Lieb JD, Myers RM, Pazin MJ, Ren B, Stamatoyannopoulos JA, Weng Z, White KP, Hardison RC. 2014. Defining functional DNA elements in the human genome. *Proceedings of the National Academy of Sciences* 111(7): 6131–6138.
- Laffin JJS, Raca G, Jackson Ca, Strand Ea, Jakielski KJ, Shriberg LD. 2012. Novel candidate genes and regions for childhood apraxia of speech identified by array comparative genomic hybridization. *Genet Med* 14 (11):928–936. doi: 10.1038/gim.2012.72
- Lai CS, Fisher SE, Hurst JA, Vargha-Khadem F, Monaco AP. 2001. A forkhead-domain gene is mutated in a severe speech and language disorder. *Nature* 413(6855):519–523. doi: 10.1038/35097076
- Lamb AN, Rosenfeld Ja, Neill NJ, Talkowski ME, Blumenthal I, Girirajan S, Shaffer LG. 2012. Haploinsufficiency of *SOX5* at 12p12.1 is associated with developmental delays with prominent language delay, behavior problems, and mild dysmorphic features. *Hum Mutat* 33(4):728–740. doi: 10.1002/humu.22037
- Lewis BA, Freebairn LA, Hansen AJ, Iyengar SK, Taylor HG. 2004. School-age follow-up of children with childhood apraxia of speech. *Lang Speech Hear Serv Sch* 35(April):122–140.
- Maunakea AK, Nagarajan RP, Bilenky M, Ballinger TJ, D'Souza C, Fouse SD, Johnson BE, Hong C, Nielson C, Zhao Y, Turecki G, Delaney A, Varhol R, Thiessen N, Shchors K, Heine VM, Rowitch DH, Xing X, Fiore C, Schillebeeckx M, Jones SJ, Haussler D, Marra MA, Hirst M, Wang T, Costello JF. 2010. Conserved role of intragenic DNA methylation in regulating alternative promoters. *Nature* 466(7303):253–257.
- Marquardt T. 2004. Token-to-token variability in developmental apraxia of speech: Three longitudinal case studies. *Clin Linguist Phon* 18(2): 127–144.
- Moreno-De-Luca A, Helmers SL, Mao H, Burns TG, Melton AMa, Schmidt KR. 2011. Adaptor protein complex-4 (AP-4) deficiency causes a novel autosomal recessive cerebral palsy syndrome with microcephaly and intellectual disability. *J Med Genet* 48(2):141–144. doi: 10.1136/jmg.2010.082263
- Peñagarikano O, Geschwind D. 2012. What does *CNTNAP2* reveal about autism spectrum disorder?. *Trends Mol Med* 18(3):156–163.
- Peter B, Raskind WH, Matsushita M, Lisowski M, Vu T, Berninger VW, Brkanac Z. 2011. Replication of *CNTNAP2* association with nonword repetition and support for *FOXP2* association with timed reading and motor activities in a dyslexia family sample. *J Neurodev Disord* 3(1):39–49. doi:10.1007/s11689-010-9065-0
- Petrin AL, Giacheti CM, Maximino LP, Abramides DVM, Zanchetta S, Rossi NF, Murray JC. 2010. Identification of a microdeletion at the 7q33-q35 disrupting the *CNTNAP2* gene in a Brazilian stuttering case. *Am J Med Genet Part A* 152A(12):3164–3172. doi: 10.1002/ajmg.a.33749
- Poliak S, Gollan L. 2001. Localization of *Caspr2* in myelinated nerves depends on axon?glia interactions and the generation of barriers along the axon The Journal of... Retrived from <http://www.jneurosci.org/content/21/19/7568.short>
- Poliak S, Gollan L, Martinez R, Custer A. 1999. *Caspr2*, a new member of the neurexin superfamily, is localized at the juxtaparanodes of myelinated axons and associates with K<sup>+</sup> channels. *Neuron* 21(19): 7568–7575.
- Poot M, Beyer V, Schwaab I, Damatova N. 2010. Disruption of *CNTNAP2* and additional structural genome changes in a boy with speech delay and autism spectrum disorder. *Neurogenetics* 11(1):81–89.
- Rasband M. 2004. It's "juxta" potassium channel! *J Neurosci Res* 76(6): 749–757.
- Reynolds CR, Kamphaus RW. 2003. RIAS, Reynolds Intellectual Assessment Scales. Psychological Assessment Resources.
- Rodenas-Cuadrado P, Ho J, Vernes SC. 2014. Shining a light on *CNTNAP2*: Complex functions to complex disorders. *Eur J Hum Genet: EJHG* 22(2): 171–178. doi: 10.1038/ejhg.2013.100
- Schafer I, Robin N, Posch J, Clark B, Izumo S, Schwartz S. 2001. Distal 5q deletion syndrome: Phenotypic correlations. *Am J Med Genet* 103(1): 63–68.
- Schneider E, Hajj N, El, Richter S. 2014. Widespread differences in cortex DNA methylation of the "language gene" *CNTNAP2* between humans and chimpanzees. *Epigenetics* 9(4):533–545.
- Semel E, Wiig E, Secord W. 2003. Clinical evaluation of language fundamentals-IV. Marickville: Harcourt Assessment.
- Shriberg LD, Aram DM, Kwiatkowski J. 1997. Developmental Apraxia of Speech. Descriptive and Theoretical Perspectives. *J Speech Lang Hear Res* 40(2): 273–285.
- Shriberg L, Tomblin J, McSweeney J. 1999. Prevalence of speech delay in 6-year-old children and comorbidity with language impairment. *J Speech Lang Hear Res* 42(6):1461–1481.
- Shriberg LD, Potter NL, Strand EA. 2011. Prevalence and phenotype of childhood apraxia of speech in youth with galactosemia. *Journal of Speech, Language, and Hearing Research* 54(2):487–519.

- Stothard SE, Snowling MJ, Bishop DVM, Chipchase BB, Kaplan CA. 1998. Language-Impaired Preschoolers. *J Speech Lang Hear Res* 41(2):407. doi: 10.1044/jslhr.4102.407
- Teperovsky EG, Bickel JO, Feldman HM. 2009. Functional characteristics of children diagnosed with childhood apraxia of speech. *Disabil Rehabil* 31(2):94–102. doi: 10.1080/09638280701795030
- Vernes SC, Newbury DF, Abrahams BS, Winchester L, Nicod J, Groszer M, Fisher SE. 2008. A functional genetic link between distinct developmental language disorders. *New Engl J Med* 359(22):2337–2345. doi: 10.1056/NEJMoa0802828
- Vogler C, Gschwind L, Röthlisberger B, Huber A, Filges I, Miny P, Papassotiropoulos A. 2010. Microarray-based maps of copy-number variant regions in European and sub-Saharan populations. *PloS One* 5(12):e15246. doi: 10.1371/journal.pone.0015246
- Woodcock RW. 1998. Woodcock reading mastery tests, revised. Circle Pines, MN: American Guidance Service.
- Worthey EA, Raca G, Laffin JJ, Wilk BM, Harris JM, Jakielski KJ, Shriberg LD. 2013. Whole-exome sequencing supports genetic heterogeneity in childhood apraxia of speech. *J Neurodevel Disord* 5(1):29.
- Zeeland AS-V. 2010. Altered functional connectivity in frontal lobe circuits is associated with variation in the autism risk gene CNTN AP2. *Sci Translat Med* 2(56):56ra80–56ra80.